Listing of claims:

The following is a complete listing of all claims in the application, with an indication of the status of each:

1. (Original) A method for lowering and controlling intraocular pressure and/or treating a mammal suffering from glaucoma, which comprises, administering to the mammal a pharmaceutically effective amount of a compound of the following formula I:

$$\begin{array}{c|c}
R^4 \\
N-R^3 \\
X \\
A \\
CH_3
\end{array}$$

I

wherein:

 $X = OH, OR^1, OCON(R^5, R^6), or OCOR^5;$

 $Y^1 = OH, OR^1, F, OCON(R^5, R^6), or OCOR^5;$

 $Y^2 = OH$, OR^1 , $OCON(R^5, R^6)$, or $OCOR^5$, with the proviso that both Y^1 and Y^2 are not OH;

 $R^1 = C_{1-3}$ alkyl;

 $R^2 = C_{1-3}$ alkyl, Cl, Br, I, CF₃, or OR¹;

 R^3 , $R^4 = H$, C_{1-3} alkyl;

 $R^5 = C_{1-6}$ alkyl; and

 $R^6 = H, C_{1-6}$ alkyl;

and pharmaceutically acceptable salts thereof.

2. (Original) The method of claim 1, wherein for the compound of formula I:

 $R^1 = methyl;$

 $R^2 = Br$, C_{1-3} alkyl; and

$$R^3$$
, $R^4 = H$.

3. (Original) The method of claim 2, wherein for the compound of formula I;

 $Y^1 = methoxy;$

 $Y^2 = OH$, methoxy; and

the α and β carbons are in the *R* configuration.

- 4. (Original) The method of claim 1, wherein the mammal is a human and the compound is administered topically.
- 5. (Original) The method of claim 1, which further comprises, administering an intraocular pressure (IOP) lowering effective amount of an IOP lowering agent selected from the group consisting of: β -blockers, carbonic anhydrase inhibitors, α 2 agonists, prostaglandin analogs, and combinations thereof.
- 6. (Original) The method of claim 5, wherein the compound of formula I and the IOP lowering agent are administered together as a single composition.
- 7. (Original) The method of claim 1, wherein the compound of formula I is selected from the group consisting of: (-)-*erythro*-(1R,2S)-1-Hydroxy-1-(4-bromo-2,5-dimethoxyphenyl)-2-aminopropane Hydrochloride; (+)-*erythro*-(1S,2R)-1-Hydroxy-1-(4-bromo-2,5-dimethoxyphenyl)-2-aminopropane Hydrochloride; (+)-*threo*-(1S, 2S)-1-Hydroxy-1-(4-bromo-2,5-dimethoxyphenyl)-2-aminopropane Hydrochloride; (-)-*erythro*-(1R,2S)-1-Methoxy-1-(4-bromo-2,5-dimethoxyphenyl)-2-aminopropane Oxalate; (+)-*erythro*-(1S,2R)-1-Methoxy-1-(4-bromo-2,5-dimethoxyphenyl)-2-aminopropane Oxalate; (+)-*threo*-(1S,2S)-1-Methoxy-1-(4-bromo-2,5-dimethoxyphenyl)-2-aminopropane Oxalate; (-)-*threo*-(1R,2R)-1-Methoxy-1-(4-bromo-2,5-dimethoxyphenyl)-2-aminopropane Oxalate; and their pharmaceutically acceptable salts.

8. (Original) The method of claim 5, wherein the compound of formula I is: (-)-threo-(1R,2R)-1-Methoxy-1-(4-bromo-2,5-dimethoxyphenyl)-2-aminopropane Oxalate and its pharmaceutically acceptable salts.

9. (Original) A compound of the following formula I:

$$\begin{array}{c}
R^4 \\
N-R^3 \\
X \\
\beta \\
CH_3
\end{array}$$

I

wherein:

 $X = OH, OR^1, OCON(R^5, R^6), or OCOR^5;$

 $Y^1 = OH, OR^1, F, OCON(R^5, R^6), or OCOR^5;$

 $Y^2 = OH$, OR^1 , $OCON(R^5, R^6)$, or $OCOR^5$, with the proviso that both Y^1 and Y^2 are not OH;

 $R^1 = C_{1-3} \text{ alkyl};$

 $R^2 = C_{1-3}$ alkyl, Cl, Br, or I with the proviso that when X = OH, R^2 is not I or methyl;

 R^3 , $R^4 = H$, C_{1-3} alkyl;

 $R^5 = C_{1-6}$ alkyl; and

 $R^6 = H, C_{1-6}$ alky;

and pharmaceutically acceptable salts thereof.

10. (Original) The compound of claim 9, wherein for formula I:

$$R^1$$
 = methyl;
 R^2 = Br, C_{1-3} alkyl; and
 R^3 , R^4 = H.

11. (Original) The compound of claim 10, wherein for formula I:

$$Y^1$$
 = methoxy;
 Y^2 = OH, methoxy; and
the α and β carbons are in the R configuration.

- 12. (Original) The compound of claim 9, which is selected from the group consisting of: (-)-(erythro-(1R,2S)-1-Hydroxy-1-(4-bromo-2,5-dimethoxyphenyl)-2-aminopropane Hydrochloride; (+)-erythro-(1S,2R)-1-Hydroxy-1-(4-bromo-2,5-dimethoxyphenyl)-2-aminopropane Hydrochloride; (-)-threo-(1R,2R)-1-Hydroxy-1-(4-bromo-2,5-dimethoxyphenyl)-2-aminopropane Hydrochloride; (-)-erythro-(1R,2R)-1-Methoxy-1-(4-bromo-2,5-dimethoxyphenyl)-2-aminopropane Oxalate; (+)-erythro-(1S,2R)-1-Methoxy-1-(4-bromo-2,5-dimethoxyphenyl)-2-aminopropane Oxalate; (+)-threo-(1S,2S)-1-Methoxy-1-(4-bromo-2,5-dimethoxyphenyl)-2-aminopropane Oxalate; (-)-threo-(1R,2R)-1-Methoxy-1-(4-bromo-2,5-dimethoxyphenyl)-2-aminopropane Oxalate; (and their pharmaceutically acceptable salts.
- 13. (Original) The compound of claim 12, which is: (-)-threo-(1R,2R)-1-Methoxy-1-(4-bromo-2,5-dimethoxyphenyl)-2-aminopropane Oxalate.